


List of things this talk will discuss:

- Kinematic Fit Cut Value Plots (new and old)
- QValue Study
- Kaon Timing Detector Study
- Unused Energy Cut
- Preliminary results of entire data set

Kin Fit Results


#Phi Events Vs Signal Confidence Level Cut

ϕ Events and ϕ Signal to Background Ratio Vs. Signal Confidence Level Cut




#Phi Events Vs BG Confidence Level Cut

ϕ Events and ϕ Signal to Background Ratio Vs. BG Confidence Level Cut



#Phi Events Vs Signal Confidence Level Cut (more points)

ϕ Events and ϕ Signal to Background Ratio Vs. Signal Confidence Level Cut



Thoughts on the Kinematic Fitter:

- CONS:
- The Kinematic Fit cut does not cut the proportional amount of signal (removes too much signal, loss of statistics)
- The fitter is a function of covariance matrices***
- PROS:
- Clearly improves signal to background ratio
- Appears to produce an exclusive Phi Eta final state
- Is there another approach or cut we can make to reduce the same amount of background while preserving more of the signal statistics?

The Qvalue Approach:


- Idea came from a paper written by Mike Williams and Curtis Meyer “Separating Signals from Non-Interfering Backgrounds using Probabilistic Event Weightings.”
- Algorithm:
 - Involves a double loop over all events
 - For a given event, calculate a kinematic distance between that event and all other events
 - Only accept the N nearest neighbors to that event
 - Plot the invariant mass that you are interested in using only the nearest neighbors
 - Fit the invariant mass distribution with a Gaussian plus a polynomial background
 - Estimate the number of signal and background events with the fits
 - Calculate the $Q\text{value} = s / (s+bg)$
- The idea is if an event close to a resonance, it will have a higher QValue than an event that is farther away

My Qvalue Study:


- I tried the study with two different kinematic distance measures:
 - One considered the ϕ distributions in the GJ frame
 - The other considered beam energy, t , missing energy, missing mass squared, and the kaon distributions in the HE frame
- Since the code is doing a double loop over the data, picking the nearest neighbors, and performing a fit; the time it took to test this approach was very long.
- Therefore, I only looked at two regions for the K^+K^- invariant mass; one that included the ϕ and one that included the ρ background.

Qvalue Example Hists

Initial Hist (All Data)




Example nearest neighbor fit




QValue Results for first kinematic distance hypothesis


Selected phi region
init_hist




Selected rho region
init_hist



QValue for phi region
sig2bgIngl_hist




QValue for rho region
sig2bgIngl_hist




QValue Results for second kinematic distance hypothesis


Selected phi region
init_hist




Selected rho region
init_hist



QValue for phi region
sig2bgIngl_hist




QValue for rho region
sig2bgIngl_hist




Invariant Mass results using QValues


phi region, first kinematic hypo




rho region, first kinematic hypo



phi region, second kinematic hypo




rho region, second kinematic hypo







Thoughts on the QValue:

- Clearly did not show any sign of being able to differentiate between signal and background for this channel.
- Even if it did work, it would have been a headache to implement
- Running over the phi or rho regions took about 3hrs + 45 min for only ~1000 events!
- Is there another approach or cut we can make to reduce the background while preserving signal statistics? (Yes, next slide)


Where does the majority of the background in the K^+K^- invariant mass spectrum come from?



Kaon Timing Plots other than TOF:




K⁺K⁻ Invariant mass Vs K⁺/⁻ Timing Detectors




K+K- Invariant mass Vs K+/- Timing Detectors Projections


ProjectionY of binx=1 [x=0.0..1.0] [$K_{TOF}^+ K_{TOF}^-$]



ProjectionY of binx=2 [x=1.0..2.0] [$K_{TOF}^+ K_{BCAL}^-$]




ProjectionY of binx=3 [x=2.0..3.0] [$K_{TOF}^+ K_{FCAL}^-$]




K+K- Invariant mass Vs K+/- Timing Detectors Projections


ProjectionY of binx=4 [x=3.0..4.0] [$K_{BCAL}^+ K_{BCAL}^-$]



ProjectionY of binx=5 [x=4.0..5.0] [$K_{BCAL}^+ K_{TOF}^-$]




ProjectionY of binx=6 [x=5.0..6.0] [$K_{BCAL}^+ K_{FCAL}^-$]




K+K- Invariant mass Vs K+/- Timing Detectors Projections


ProjectionY of binx=7 [x=6.0..7.0] [K_{FCAL}^+ K_{FCAL}^-]



ProjectionY of binx=8 [x=7.0..8.0] [K_{FCAL}^+ K_{TOF}^-]



ProjectionY of binx=9 [x=8.0..9.0] [K_{FCAL}^+ K_{BCAL}^-]




Thoughts on the Kaon Timing:

- The best way to identify the phi with my channel is using only the TOF and nothing else
- There may be other detectors that observe a phi. However, there appears to be so much background associated with the FCAL/BCAL that it seems more logical to just throw them out
- Initially I didn't want to perform a cut like this out of fear that it would hurt any possibility of performing an angular distribution study
- You both may think of a better way to perform this cut. But for now I allow kaon timing to only come from the TOF.

Kaon Timing Cut Impact


Before Cut



Fit Results:

Signal Events: 533
 Background Events: 518
 S/BG: 1.029
 Total BG: 25417

After Cut




Fit Results:


Signal Events: 469 (-12%)
 Background Events: 219 (-58%)
 S/BG: 2.138 (+208%)
 Total BG: 3293 (-87%)

Unused Energy Cut (after TOF Cut)

Unused Energy Distribution




K+ K- Mass Vs Unused Energy Distribution




K+ K- Mass Projection; Unused Energy = 0

ProjectionY of binx=1 [x=0.000..0.008]




K+ K- Mass Projection; Unused Energy > 0

ProjectionY of binx=[12,511] [x=0.092..4.258]



TOF+Unused Energy Cut Impact


Before Cuts



Fit Results:

Signal Events: 533
 Background Events: 518
 S/BG: 1.029
 Total BG: 25417

After Cuts



Fit Results:


Signal Events: 303 (-43%)
 Background Events: 94 (-82%)
 S/BG: 3.205 (+311%)
 Total BG: 1244 (-95%)

Preliminary List of Final Cuts:


- Delta T for each particle species and sub detector
 - Beam Energy Cut
 - Beam Bunch Cut (RF Time)
 - Vertex Cuts
 - P vs Theta Cut for Photons (Reduces Secondaries)
 - Number of photons reconstructed in the event
 - Kaons are only allowed to get timing from TOF
 - Unused Energy < 50 MeV
-
- NOT INCLUDED: Signal Kinematic Fitter Confidence Level
 - NOT INCLUDED: Background Kinematic Fitter Confidence Level

Results for All Data, All Cuts


K^+K^- Invariant Mass (GeV/c^2) [2σ]




$\gamma\gamma$ Invariant Mass (GeV/c^2) [2σ]







Missing Mass Squared (GeV/c^2)² [2σ]



Missing Energy (GeV) [2σ]




Results for All Data, All Cuts




Results for All Data, All Cuts


Mandelstam t' - Large $(\text{GeV}/c^2)^2$ [2σ]




Mandelstam t' - Signal+Background $(\text{GeV}/c^2)^2$ [2σ]







Mandelstam t' - Background $(\text{GeV}/c^2)^2$ [2σ]







Mandelstam t' - Signal $(\text{GeV}/c^2)^2$ [2σ]







Results for All Data, All Cuts



Results for All Data, All Cuts




Results for All Data, All Cuts




Results for All Data, All Cuts


pK⁻ Invariant Mass - Large Sample (GeV/c²) [2 σ]




pK⁻ Invariant Mass - Signal+Background (GeV/c²) [2 σ]







pK⁻ Invariant Mass - Background (GeV/c²) [2 σ]







pK⁻ Invariant Mass - Signal (GeV/c²) [2 σ]



Results for All Data, All Cuts



Results for All Data, All Cuts



Results for All Data, All Cuts

