

Baryon Spectroscopy at ELSA / Jefferson Lab

What have we learned about excited baryons?

Volker Credé

Florida State University, Tallahassee, FL

Physics Colloquium

University of Bonn, 11/30/2012



Outline

- 1 Introduction
 - Quarks, QCD, and Confinement
 - Structure of Baryon Resonances
- 2 The Search for Undiscovered States
 - Electromagnetic Probes
 - Mission Goal: Complete Experiments
- 3 Results from Photoproduction Experiments
 - Photoproduction of π , η , and ω Mesons
 - Observables in the Photoproduction of Two Pions
- 4 Summary and Outlook



Outline

1

Introduction

- Quarks, QCD, and Confinement
- Structure of Baryon Resonances

2

The Search for Undiscovered States

- Electromagnetic Probes
- Mission Goal: Complete Experiments

3

Results from Photoproduction Experiments


- Photoproduction of π , η , and ω Mesons
- Observables in the Photoproduction of Two Pions

4

Summary and Outlook



Non-Perturbative QCD



QCD is the theory of the strong nuclear force which describes the interactions of quarks and gluons making up hadrons.

Strong processes at larger distances and at small (soft) momentum transfers belong to the realm of non-perturbative QCD.


Quarks are confined within hadrons.

Confinement of quarks and gluons within nucleons is a non-perturbative phenomenon, and QCD is extremely hard to solve in non-perturbative regimes: Knowledge of internal structure of nucleons is still limited.

This is particularly true for excited nucleons.



Non-Perturbative QCD




How does QCD give rise to excited hadrons?


Interaction between quarks fairly unknown
throughout $> 98\%$ of a hadron's volume.

Explaining the excitation spectrum of hadrons is central to our understanding of QCD in the low-energy regime (Hadron Models, Lattice QCD, etc.)

→ Complementary to Deep Inelastic Scattering (DIS) where information on collective degrees of freedom is lost.




Non-Perturbative QCD



How does QCD give rise to excited hadrons?

- 1 What is the origin of confinement?
- 2 How are confinement and chiral symmetry breaking connected?
- 3 Would the answers to these questions explain the origin of $\sim 98\%$ of observed matter in the universe?

Excited Baryons: What are the fundamental degrees of freedom inside a proton or a neutron? How do they change with varying quark masses?



CQM



CQM+flux tubes




Quark-diquark clustering



Nucleon-meson system

Understanding Excited Baryons

The excitation spectrum of the hydrogen atom:





Study of the hydrogen spectrum has shown that the understanding of the structure of a bound state and of its excitation spectrum go hand in hand.

Understanding Excited Baryons

	****	***	**	*
N^*	11	3 → 5	6 → 8	2 → 3
Δ^*	7	3	6 → 7	6 → 5

PDG 2012




Nomenclature:

 $P_{33}(1232)$ (old) $\Delta(1232) \frac{3}{2}^{+}$ (new)


Spectrum of Nucleon Resonances

— S. Capstick and N. Isgur, Phys. Rev. **D34** (1986) 2809




Spectrum of Nucleon Resonances

— S. Capstick and N. Isgur, Phys. Rev. **D34** (1986) 2809



Excited-State Baryon Spectroscopy from Lattice QCD

R. Edwards *et al.*, Phys. Rev. D **84**, 074508 (2011)



Exhibits broad features expected of $SU(6) \otimes O(3)$ symmetry

→ Counting of levels consistent with non-rel. quark model, no parity doubling

Components of the Experimental N^* Program

The excited baryon program has two main components:

- **Probe resonance transitions at different distance scales**

Electron beams are ideal to measure resonance form factors and their corresponding Q^2 dependence.



→ Provides information on the structure of excited nucleons and on the confining (effective) forces of the 3-quark system.

- **Establish the systematics of the spectrum**

Current medium-energy experiments use photon beams to map out the baryon spectrum (JLab, ELSA, MAMI, SPring-8, etc.).

→ Provides information on the nature of the effective degrees of freedom in strong QCD and also addresses the issue of previously unobserved or so-called *missing resonances*.

Helicity Amplitudes for the “Roper” Resonance



Data from CLAS

$A_{1/2}$ and $S_{1/2}$ amplitudes:

e.g. I. Aznauryan *et al.*,
PRC **78**, 045209 (2008);
PRC **80**, 055203 (2009).

Quark-model calculations:

— q^3G hybrid state



--- q^3 radial excitation

Consistency between both channels ($N\pi\pi$, $N\pi$): sign change, magnitude, ...

- At short distances (high Q^2), Roper behaves like radial excitation.
- Low Q^2 behavior not well described by LF quark models:
e.g. meson-baryon interactions missing

→ Gluonic excitation likely ruled out!


Helicity Amplitudes for $\gamma^* p \rightarrow N(1520)D_{13}$ Transition



There is clear evidence for helicity switch from $\lambda = 3/2$ (at photon point) to $\lambda = 1/2$ at high Q^2 :

- Rapid change in helicity structure when going from photo- to electroproduction of a nucleon resonance
→ Stringent prediction of the CQM!

$$A_{\text{hel}} = \frac{|A_{1/2}|^2 - |A_{3/2}|^2}{|A_{1/2}|^2 + |A_{3/2}|^2} \rightarrow$$



Components of the Experimental N^* Program

The excited baryon program has two main components:

- **Probe resonance transitions at different distance scales**

Electron beams are ideal to measure resonance form factors and their corresponding Q^2 dependence.

→ Provides information on the structure of excited nucleons and on the confining (effective) forces of the 3-quark system.

- **Establish the systematics of the spectrum**


Current medium-energy experiments use photon beams to map out the baryon spectrum (JLab, ELSA, MAMI, SPring-8, etc.).

→ Provides information on the nature of the effective degrees of freedom in strong QCD and also addresses the issue of previously unobserved or so-called *missing resonances*.

Outline


- 1 Introduction
 - Quarks, QCD, and Confinement
 - Structure of Baryon Resonances
- 2 The Search for Undiscovered States
 - Electromagnetic Probes
 - Mission Goal: Complete Experiments
- 3 Results from Photoproduction Experiments
 - Photoproduction of π , η , and ω Mesons
 - Observables in the Photoproduction of Two Pions
- 4 Summary and Outlook





Experimental Facilities

CBELSA/TAPS at ELSA



Meson photoproduction:


- $\gamma p \rightarrow p \pi^0 \rightarrow p \gamma \gamma$
- $\gamma p \rightarrow p \eta \rightarrow p \gamma \gamma, p 3\pi^0, p \pi^+ \pi^- \pi^0$
- $\gamma p \rightarrow p \omega \rightarrow p \pi^0 \gamma, \pi^+ \pi^- \pi^0$

Jefferson Laboratory




Double-Polarization: Toward Complete Experiments

Calorimeter system at ELSA is optimized for neutral particles.



Crystal Barrel




Close to 4π coverage

Crystal Barrel Setup at ELSA

- Crystal Barrel (1230 CsI crystals)
- Inner Detector (513 scintillating fibers)
- Forward Plug (90 CsI crystals with PM's)
- MiniTAPS (216 BaF₂, 1° - 12°)



Frozen-Spin Target: Butanol (C₄H₉OH).

The CLAS Spectrometer at Jefferson Laboratory




g8b

linear beam
polarization



FROST



double polarization



Double-Polarization: Frozen Spin Targets

Horizontal cryostat with integrated solenoid to freeze the proton spin.

- DNP at high B-field (2.5 T), holding mode at 0.4 T
- Relaxation time at ELSA ~ 500 h



“ELSA”



“CLAS”

Transverse Target Polarization
(race-track coil - Dipole Magnet)


Longitudinally-Polarized Target ($P_z \approx 80\%$)

Extraction of Resonance Parameters

- Double-polarization measurements
- Measurements off neutron and proton to resolve isospin contributions:
 - 1 $\mathcal{A}(\gamma N \rightarrow \pi, \eta, K)^{I=3/2} \iff \Delta^*$
 - 2 $\mathcal{A}(\gamma N \rightarrow \pi, \eta, K)^{I=1/2} \iff N^*$
- Re-scattering effects: Large number of measurements (and reaction channels) needed to extract full scattering amplitude.



Coupled Channels



Why are Polarization Observables Important?

- ① From π threshold up to $\Delta(1232)$ region
 - s - & p -wave approximation
 - Fermi-Watson Theorem
 - Two observables sufficient, e.g. $d\sigma/d\Omega$, Σ .
- ② Above the $\pi\pi$ threshold
 - More observables needed.

$$\frac{d\sigma}{d\Omega} = \sigma_0 \{ 1 - \delta_I \Sigma \cos 2\phi + \Lambda_x (-\delta_I H \sin 2\phi + \delta_\odot F) - \Lambda_y (-T + \delta_I P \cos 2\phi) - \Lambda_z (-\delta_I G \sin 2\phi + \delta_\odot E) \}$$

Chiang & Tabakin, Phys. Rev. C **55**, 2054 (1997)

In order to determine the full scattering amplitude without ambiguities, one has to carry out eight carefully selected measurements: four double-spin observables along with four single-spin observables.

Eight well-chosen measurements are needed to fully determine production amplitudes F_1 , F_2 , F_3 , and F_4 .

Why are Polarization Observables Important?

- 1 From π threshold up to $\Delta(1232)$ region
 - s- & p-wave approximation
 - Fermi-Watson Theorem
 - Two observables sufficient, e.g. $d\sigma/d\Omega$, Σ .

$$\begin{aligned} \frac{d\sigma}{d\Omega} = \sigma_0 \{ & 1 - \delta_I \Sigma \cos 2\phi \\ & + \Lambda_x (-\delta_I H \sin 2\phi + \delta_\odot F) \\ & - \Lambda_y (-T + \delta_I P \cos 2\phi) \\ & - \Lambda_z (-\delta_I G \sin 2\phi + \delta_\odot E) \} \end{aligned}$$

- 2 Above the $\pi\pi$ threshold
 - More observables needed.

Chiang & Tabakin, Phys. Rev. C55, 2054 (1997)

In order to determine the full scattering amplitude without ambiguities, one has to carry out eight carefully selected measurements: four double-spin observables along with four single-spin observables.


Eight well-chosen measurements are needed to fully determine production amplitudes F_1 , F_2 , F_3 , and F_4 .

Example: Ambiguities in $\gamma p \rightarrow p \pi^0$

— Bonn-Gatchina (BnGa)

— SAID

— MAID



$$\frac{d\sigma}{d\Omega} = \sigma_0 \{ 1 - \delta_I \Sigma \cos 2\phi$$

$$+ \Lambda_x (-\delta_I H \sin 2\phi + \delta_\odot F)$$

$$- \Lambda_y (-T + \delta_I P \cos 2\phi)$$

$$- \Lambda_z (-\delta_I G \sin 2\phi + \delta_\odot E) \}$$




Total cross section: $\gamma p \rightarrow p \pi^0$

Example: Ambiguities in $\gamma p \rightarrow p \pi^0$


— Bonn-Gatchina (BnGa) PWA

— SAID

— MAID



Helicity Asymmetry E in $\vec{\gamma} \vec{p} \rightarrow p \pi^0$ (Data from ELSA)



$$E = \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}}$$


$$E_\gamma \in [0.6, 2.2] \text{ GeV}$$

- CBELSA/TAPS
- Maid
- Said
- BoGa

Angular distributions
sensitive to interference
between resonances.

M. Gottschall *et al.* [CBELSA/TAPS]



$\cos \theta_{\pi^0}$

Example: Ambiguities in $\gamma p \rightarrow p \eta$ 

$$\frac{d\sigma}{d\Omega} = \sigma_0 \{ 1 - \delta_I \Sigma \cos 2\phi \}$$


CLAS g11a ■
CBELSA/TAPS ○
MAMI ●
GRAAL ▼

BoGa-PWA

 $E_\gamma = 1250 \text{ MeV}$  η -MAIDD. Elsner *et al.* [CBELSA/TAPS], EPJ A **33**, 147 (2007)



Helicity-Dependent Cross Section for $\vec{\gamma} \vec{p} \rightarrow p \eta$

η count rate difference



$$E = \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}}$$

→ Very preliminary:
Data are positive






M. Gottschall et al. [CBELSA/TAPS]

B. Morrison et al. [CLAS Collaboration]

Systematics in Photoproduction (Normalization)

$$\frac{d\sigma}{d\Omega} = \frac{N_{\text{events}}}{\Phi_{\gamma} \epsilon_{\text{MC}} \rho_t \Gamma \Delta\Omega}$$




$\gamma p \rightarrow p\eta$: O. Bartholomy, V.C. *et al.* [ELSA]

PRL **94** (2005), EPJ A **33** (2007), PRC **80** (2009).

Outline

- 1 Introduction
 - Quarks, QCD, and Confinement
 - Structure of Baryon Resonances
- 2 The Search for Undiscovered States
 - Electromagnetic Probes
 - Mission Goal: Complete Experiments
- 3 Results from Photoproduction Experiments
 - Photoproduction of π , η , and ω Mesons
 - Observables in the Photoproduction of Two Pions
- 4 Summary and Outlook



Isospin Filter: $\gamma p \rightarrow N^* (I = 1/2) \rightarrow p \omega$ A. Wilson *et al.* [CBELSA/TAPS], Ph.D. thesis (Florida State)


— CBELSA/TAPS

-- CLAS (2009)

Good agreement
between experiments

Excellent statistics

Great progress in our
understanding of the
differences between
experiments.

Isospin Filter: $\gamma p \rightarrow N^* (I = 1/2) \rightarrow p \omega$ A. Wilson *et al.* [CBELSA/TAPS], Ph.D. thesis (Florida State)


— CBELSA/TAPS

— CLAS (2009)

Good agreement
between experiments


Excellent statistics

New CBELSA/TAPS
data extend the world
database in the most
forward direction.

Isospin Filter: $\gamma p \rightarrow N^* (I = 1/2) \rightarrow p \omega$ A. Wilson *et al.* [CBELSA/TAPS], Ph.D. thesis (Florida State)



— CBELSA/TAPS

— CLAS (2009)



Baryon Resonances in the Reaction $\gamma p \rightarrow p \omega$

M. Williams *et al.* [CLAS Collaboration], Phys. Rev. C **80**, 065209 (2009)




PWA fit includes resonances + t -channel amplitudes.


Strong evidence for ($W > 2 \text{ GeV}$):

$(5/2)+ N(1680)$ * * *

$(5/2)+ N(1950)$ **

$(7/2)- N(2190)$ * * *




Polarization Observables in $\gamma p \rightarrow p \omega$ 

Strong evidence for ($W > 2$ GeV):

($5/2+$) + N(1680) * * *

($5/2+$) + N(1950) * *

($7/2-$) - N(2190) * * *

Asymmetry Σ for $\gamma p \rightarrow p \omega$ (P. Collins *et al.*, CUA)

— Oh *et al.*


— Paris *et al.*

— Sarantsev *et al.*

Photoproduction of π^0 Mesons from the Proton

Reaction $\gamma p \rightarrow p \pi^0$ remains important for our understanding of baryons.


- At ELSA, excellent data with good statistics in the forward direction.
- Forward region is very sensitive to higher-spin resonances:
 → Observation of $N(2190)G_{17}$ within the Bonn-Gatchina PWA framework
 (Important to confirm high-mass states first observed in πN scattering)



V. C. *et al.* [CBELSA/TAPS], Phys. Rev. C **84**, 055203 (2011)

Beam Asymmetry Σ in $\vec{\gamma} p \rightarrow p \pi^0$

$$\gamma p \rightarrow p \pi^0$$



$$\frac{d\sigma}{d\Omega} = \sigma_0 \{ 1 - \delta_I \Sigma \cos 2\phi$$


$$+ \Lambda_x (-\delta_I H \sin 2\phi + \delta_\odot F) \\ - \Lambda_y (-T + \delta_I P \cos 2\phi) \\ - \Lambda_z (-\delta_I G \sin 2\phi + \delta_\odot E) \}$$

— SAID — MAID ● CLAS
($E_\gamma < 2$ GeV, $-0.85 < \cos \theta_\pi < -0.35$)

→ Serious discrepancies between models and data above 1.4 GeV.

Photoproduction of π mesons still not very well understood.

M. Dugger (ASU), CLAS g8b run group, to be published

Beam Asymmetry Σ in $\vec{\gamma} p \rightarrow p \pi^0$ 

Combination of $p \pi^0$ and $n \pi^+$ final states can help distinguish between Δ and N^* resonances:



$$\begin{array}{cc}
 \Delta^+ & N^* \\
 \downarrow & \downarrow \\
 \pi^0 + p : \sqrt{2/3} \left| I = \frac{3}{2}, I_3 = \frac{1}{2} \right\rangle - \sqrt{1/3} \left| I = \frac{1}{2}, I_3 = \frac{1}{2} \right\rangle & \\
 \pi^+ + n : \sqrt{1/3} \left| I = \frac{3}{2}, I_3 = \frac{1}{2} \right\rangle + \sqrt{2/3} \left| I = \frac{1}{2}, I_3 = \frac{1}{2} \right\rangle &
 \end{array}$$

M. Dugger (ASU), CLAS g8b run group, to be published

Beam Asymmetry Σ in $\vec{\gamma}p \rightarrow p \pi^0$ and $\vec{\gamma}p \rightarrow n \pi^+$

$$\gamma p \rightarrow p \pi^0$$

$$\gamma p \rightarrow \pi^+ n$$




M. Dugger (ASU), CLAS g8b run group, to be published

Asymmetry G in $\vec{\gamma} \vec{p} \rightarrow p \pi^0$ (Results from ELSA)

$$\frac{d\sigma}{d\Omega} = \sigma_0 \{ 1 - \delta_I \Sigma \cos 2\phi$$


$$+ \Lambda_x (-\delta_I H \sin 2\phi + \delta_\odot F) \\ - \Lambda_y (-T + \delta_I P \cos 2\phi) \\ - \Lambda_z (-\delta_I G \sin 2\phi + \delta_\odot E) \}$$



Surprisingly, π production also not well understood at lower energies:

— BoGa
-- SAID
... MAID

A. Thiel *et al.* [CBELSA/TAPS], PRL **109**, 102001 (2012)

Asymmetry G in $\vec{\gamma} \vec{p} \rightarrow p \pi^0$ (Results from ELSA)

$$\frac{d\sigma}{d\Omega} = \sigma_0 \{ 1 - \delta_I \Sigma \cos 2\phi$$

$$+ \Lambda_x (-\delta_I \textcolor{blue}{H} \sin 2\phi + \delta_{\odot} \textcolor{blue}{F}) \\ - \Lambda_y (-\textcolor{blue}{T} + \delta_I \textcolor{blue}{P} \cos 2\phi) \\ - \Lambda_z (-\delta_I \textcolor{red}{G} \sin 2\phi + \delta_{\odot} \textcolor{blue}{E}) \}$$

$$\theta_{\pi} = 90 \pm 5^{\circ}$$

Surprisingly, π production also not well understood at lower energies.

$$\theta_{\pi} = 130 \pm 5^{\circ}$$


Below 1 GeV, discrepancies can be traced to the E_{0+} and E_{2-} multipoles, which are related to certain resonances:

$$E_{0+}: N(1535) \frac{1}{2}^{-}, N(1650) \frac{1}{2}^{-}, \Delta(1620) \frac{1}{2}^{-}$$


$$E_{2-}: N(1520) \frac{3}{2}^{-}, \Delta(1700) \frac{3}{2}^{-}$$

Transverse Target Polarization: Target Asymmetry T

$700 < E_\gamma < 800$ MeV



$800 < E_\gamma < 900$ MeV



direction of
target pol.:


$\beta = 99^\circ$

$$\Delta N(\phi) = \frac{1}{f P_{\text{target}}} \cdot \frac{N_{\uparrow} - N_{\downarrow}}{N_{\uparrow} + N_{\downarrow}}$$

$$= T \cdot \sin(\phi - \beta)$$

→ Unprecedented statistical quality.

photon pol.		target pol. axis		
		x	y	z
unpolarized	σ		T	
linear	$-\Sigma$	H	$-P$	$-G$
circular		F		$-E$

Target Asymmetry T in $\gamma \vec{p} \rightarrow p \pi^0$ (Data from ELSA)

$600 < E_\gamma < 1000$ MeV
in bins of $\Delta E = 25$ MeV

● CBELSA/TAPS

▼ Daresbury (1977)

Nucl. Phys. B **121** (1977), 45


— Maid

— Said

— Bonn Gatchina

➔ Good agreement of
experimental data.
(improved statistics)

J. Hartmann *et al.* [CBELSA/TAPS]

Target Asymmetry T in $\gamma \bar{p} \rightarrow p \pi^0$ (Data from ELSA) $1000 < E_\gamma < 1500 \text{ MeV}$

● CBELSA/TAPS

▼ Daresbury (1977)


Nucl. Phys. B **121** (1977), 45

— Maid

— Said

— Bonn Gatchina

→ Good agreement of
experimental data.Model disagreement
toward 3rd resonance
region: $W > 1.6 \text{ GeV}$
(E_{0+}, E_{2-} multipoles?)

Target Asymmetry T in $\gamma \bar{p} \rightarrow p \eta$ (Data from ELSA)

$708 < E_\gamma < 933$ MeV
in bins of $\Delta E = 25$ MeV

● CBELSA/TAPS

▲ PHOENICS (1998)



Phys. Rev. Lett. **81** (1998), 534

— Maid

— Said

— Bonn Gatchina

J. Hartmann *et al.* [CBELSA/TAPS]

Observables P, H (Results from CBELSA/TAPS) $800 < E_\gamma < 900$ MeVangle of lin.
pol. plane:

$\alpha = 45^\circ$

direction of
target pol.:

$\beta = 99^\circ$

$$\Delta N(\phi) = C \cdot \frac{(N_{\perp\uparrow} - N_{\perp\downarrow}) - (N_{\parallel\uparrow} - N_{\parallel\downarrow})}{(N_{\perp\uparrow} + N_{\perp\downarrow}) + (N_{\parallel\uparrow} + N_{\parallel\downarrow})}$$

$$= P (\sin(\phi - \beta) \cos(2(\phi - \alpha)))$$

$$+ H (\cos(\phi - \beta) \sin(2(\phi - \alpha)))$$


photon pol.		target pol. axis		
		x	y	z
unpolarized	σ		T	
linear	$-\Sigma$	H	$-P$	$-G$
circular		F		$-E$

Beam-Target Polarization Observables in $\gamma p \rightarrow p \pi\pi$


$$\begin{aligned}
 I = I_0 \{ & (1 + \vec{\Lambda}_i \cdot \vec{P}) \\
 & + \delta_{\odot} (I^{\odot} + \vec{\Lambda}_i \cdot \vec{P}^{\odot}) \\
 & + \delta_I [\sin 2\beta (I^S + \vec{\Lambda}_i \cdot \vec{P}^S) + \\
 & \cos 2\beta (I^C + \vec{\Lambda}_i \cdot \vec{P}^C)] \}
 \end{aligned}$$

W. Roberts *et al.*, Phys. Rev. C **71**, 055201 (2005)

⇐ Double-Meson
Final States
(15 Observables)



At higher excitation energies:
Multi-meson final states important.



Photoproduction of $\pi\pi$ Pairs off the Proton: Kinematics

Two mesons in the final state require 5 independent variables!


For example: E_γ , $\Theta_{\text{c.m.}}$, ϕ^* , θ^* , $M_{p+\text{meson}_1}$

Single-meson reactions:


→ p-meson system in the reaction plane

Two-meson reactions:

→ Reaction and decay plane
form angle ϕ



Beam Asymmetries I^S, I^C in $\vec{\gamma} p \rightarrow p \pi^0 \pi^0$




First measurements of beam asymmetries I^S and I^C using linearly-polarized photons in the reaction $\gamma p \rightarrow p \pi^0 \pi^0$.

Among other things, study of decays into $\Delta\pi$:

- BoGa-PWA solution with a dominant $\Delta(1700)D_{33} \rightarrow \Delta\pi$ D -wave
- BoGa-PWA solution with a dominant $\Delta(1700)D_{33} \rightarrow \Delta\pi$ S -wave


- Direct measurements
- From mirror operation $I^S(\Phi^*) \rightarrow I^S(2\pi - \Phi^*)$

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1100 < E_\gamma < 1150$ MeVC. Hanretty *et al.*, CLAS-g8b run group, under review

Beam Asymmetry $I^\odot$ in $\vec{\gamma}p \rightarrow p\pi^+\pi^-$ from JLab

Analysis of butanol target challenging:

- Determination of dilution factor $\rightarrow$
(event-based dilution factors possible)



S. Park *et al.*, CLAS (FROST); CLAS (g1c) PRL **95** (2005)

Outline

- 1 Introduction
 - Quarks, QCD, and Confinement
 - Structure of Baryon Resonances
- 2 The Search for Undiscovered States
 - Electromagnetic Probes
 - Mission Goal: Complete Experiments
- 3 Results from Photoproduction Experiments
 - Photoproduction of π , η , and ω Mesons
 - Observables in the Photoproduction of Two Pions
- 4 Summary and Outlook



Summary and Outlook

Our understanding of baryon resonances has made great leaps forward. There is good evidence that most of the known states (listed in the PDG) will be confirmed in photoproduction and that new states will be revealed:

- Goal of performing (almost) complete experiments has been (almost) achieved; program on neutron ongoing.



$N(1860)$	$\frac{5}{2}^{+}$	**	πN	γN				
$N(1875)$	$\frac{1}{2}^{-}$	* * *	πN	γN	ΛK	ΣK		$N\sigma$
$N(1880)$	$\frac{1}{2}^{+}$	**	πN	γN	ΛK	ΣK		
$N(1895)$	$\frac{1}{2}^{-}$	**	πN	γN	ηN	ΛK	ΣK	
$N(1900)$	$\frac{3}{2}^{+}$	* * *	πN	γN	ηN	ΛK	ΣK	$\Delta\pi$
$N(2060)$	$\frac{3}{2}^{-}$	**	πN	γN	ηN	ΣK		
$\Delta(1940)$	$\frac{3}{2}^{-}$	* $\rightarrow$ **	πN	γN				$\Delta\eta (!)$

New States
in PDG 2012.

Summary and Outlook

Our understanding of baryon resonances has made great leaps forward. There is good evidence that most of the known states (listed in the PDG) will be confirmed in photoproduction and that new states will be revealed:


- Goal of performing (almost) complete experiments has been (almost) achieved; program on neutron ongoing.
- Spectroscopy will continue at ELSA, MAMI and JLab in the 12 GeV era (e.g. Ξ , Ω states): GlueX and CLAS12.



$N(1860)_{\frac{1}{2}^{+}}$	**	πN	γN				
$N(1875)_{\frac{1}{2}^{-}}$	* * *	πN	γN	ΛK	ΣK		$N\sigma$
$N(1880)_{\frac{1}{2}^{+}}$	**	πN	γN	ΛK	ΣK		
$N(1895)_{\frac{1}{2}^{-}}$	**	πN	γN	ηN	ΛK	ΣK	
$N(1900)_{\frac{3}{2}^{+}}$	* * *	πN	γN	ηN	ΛK	ΣK	$\Delta\pi$
$N(2060)_{\frac{3}{2}^{-}}$	**	πN	γN	ηN		ΣK	
$\Delta(1940)_{\frac{3}{2}^{-}}$	* $\rightarrow$ **	πN	γN				$\Delta\eta (!)$


New States
in PDG 2012.

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1100 < E_\gamma < 1150$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1150 < E_\gamma < 1200$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1200 < E_\gamma < 1250$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1250 < E_\gamma < 1300$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1300 < E_\gamma < 1350$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1350 < E_\gamma < 1400$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1400 < E_\gamma < 1450$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1450 < E_\gamma < 1500$ MeV


C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1500 < E_\gamma < 1550$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1550 < E_\gamma < 1600$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1600 < E_\gamma < 1650$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1650 < E_\gamma < 1700$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1700 < E_\gamma < 1750$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1750 < E_\gamma < 1800$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1800 < E_\gamma < 1850$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1850 < E_\gamma < 1900$ MeV




C. Hanretty et al., CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1900 < E_\gamma < 1950$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $1950 < E_\gamma < 2000$ MeV




C. Hanretty *et al.*, CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $2000 < E_\gamma < 2050$ MeV




C. Hanretty et al., CLAS-g8b run group, under review

I^S in $\vec{\gamma} p \rightarrow p \pi^+ \pi^-$ from JLab: $2050 < E_\gamma < 2100$ MeV



C. Hanretty et al., CLAS-g8b run group, under review

Fascinating Discovery in Reactions off the Neutron



Neutron measurements important:

- Different resonance contributions
- Study of isospin composition of el.-magn. couplings

Narrow structure in $\gamma d \rightarrow n \eta + p$
with $M \approx 1670$ MeV and $\sigma = 25$ MeV:

- Interference effect of the $S_{11}(1535)$ and $S_{11}(1535)$ resonances
- Coupled channel effect of $S_{11}(1535)$ and $P_{11}(1710)$
- New narrow state?